

## TPO Human, HEK

**Description:**Thrombopoietin Human Recombinant produced in HEK cells is a glycosylated monomer, having a molecular weight range of 80-85kDa due to glycosylation.The TPO is purified by proprietary chromatographic techniques.

Catalog #:CYPs-122

**Synonyms:**Megakaryocyte colony-stimulating factor, Myeloproliferative leukemia virus oncogene ligand, C-mpl ligand, ML, Megakaryocyte growth and development factor, MGDF, TPO, MKCSF, MPLLg, MGC163194, THPO.

For research use only.

**Source:**HEK.

**Physical Appearance:**Sterile Filtered White lyophilized (freeze-dried) powder.

**Purity:**Greater than 90% as observed by SDS-PAGE.

**Formulation:**

The TPO protein was lyophilized from 1mg/ml in 1xPBS.

**Stability:**

Lyophilized Thrombopoietin although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution TPO should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.

**Usage:**

NeoBiolabs products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Solubility:**

It is recommended to reconstitute the lyophilized Thrombopoietin in sterile PBS containing 0.1% endotoxin-free recombinant HSA.

**Introduction:**

Thrombopoietin is a glycoprotein hormone produced mainly by the liver and the kidney that regulates the production of platelets by the bone marrow. It stimulates the production and differentiation of megakaryocytes, the bone marrow cells that fragment into large numbers of platelets.

**Biological Activity:**

The specific activity was determined by the dose-dependent stimulation of the proliferation of human TF-1 cells (human erythroleukemic indicator cell line) and is typically 0.5-5ng/ml.

**To place an order, please [Click HERE](#).**