

## IGF1 Human Des1-3

**Description:**IGF-I Des(1-3) Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 67 amino acids (aa 4-70) and having a molecular mass of 7372 Dalton. IGF-1 Des1-3 is purified by proprietary chromatographic techniques.

**Synonyms:**Somatomedin C, IGF-I, IGFI, IGF1, IGF-IA, Mechano growth factor, MGF, Des(1-3), Des1-3, Des 1-3, Des (1-3), IGF-1 (4-70).

**Source:**Escherichia Coli.

**Physical Appearance:**Sterile Filtered White lyophilized (freeze-dried) powder.

**Amino Acid Sequence:**The sequence of the first five N-terminal amino acids was determined and was found to be Thr-Leu-Cys-Gly-Ala.

**Purity:**Greater than 95.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

**Formulation:**

The protein was lyophilized after extensive dialysis against 50mM acetic acid buffer.

**Stability:**

Lyophilized IGF-I des(1-3) although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IGF1 des-1-3 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.

**Usage:**

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Solubility:**

It is recommended to reconstitute the lyophilized IGF-I Des(1-3) in 100mM Acetic acid not less than 100g/ml, which can then be further diluted to other aqueous solutions.


**Introduction:**

The somatomedins, or insulin-like growth factors (IGFs), comprise a family of peptides that play important roles in mammalian growth and development. IGF1 mediates many of the growth-promoting effects of growth hormone (GH; MIM 139250). Early studies showed that growth hormone did not directly stimulate the incorporation of sulfate into cartilage, but rather acted through a serum factor, termed 'sulfation factor,' which later became known as 'somatomedin' (Daughaday et al., 1972). Three main somatomedins have been characterized: somatomedin C (IGF1), somatomedin A (IGF2; MIM 147470), and somatomedin B (MIM 193190) (Rotwein, 1986; Rosenfeld, 2003).

**Biological Activity:**

The ED50, calculated by the dose-dependant proliferation of murine BALB\3T3 cells (measured by 3H-thymidine uptake) is < 1ng/ml, corresponding to a specific activity of 1,000,000units/mg. For most in-vitro applications, IGF-I des1-3 exerts its biological activity in the concentration range

www.neobiolab.com  
info@neobiolab.com  
888.754.5670, +1 617.500.7103 United States  
0800.088.5164, +44 020.8123.1558 United Kingdom  
of 0.2-20 ng/ml, corresponding to a specific activity of 50,000-5,000,000units/mg.



***To place an order, please [Click HERE](#).***

Catalog #:CYP5-525

For research use only.

© 2015 NeoScientific



Request a  
CATALOG  
[CLICK HERE](#)