www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

PKC-g Human

Description: PKC-g Human Recombinant produced in Sf9 is a glycosylated, polypeptide chain containing amino acids 2-676 and having a molecular mass of 80 KD. This protein is the full-length form of the protein with an amino terminal poly His-tag.PKC-g is purified by proprietary chromatographic techniques.

Catalog #:PKPS-327

For research use only.

Synonyms: Protein kinase C gamma type, EC 2.7.11.13, PKC-gamma, PRKCG, PKCC, PKCG, SCA14, MGC57564.

Source:Sf9 insect cells.

Physical Appearance: Sterile Filtered clear solution.

Purity: Greater than 90% as determined by SDS-PAGE.

Formulation:

PKC-g is supplied at a of 0.1mg/ml in 10mM Tris, pH 7.4, 0.1M NaCl, 20% glycerol, 1mM DTT, 0.1mM EDTA, 0.2mM PMSF and 0.03% Brij-35.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play distinct roles in cells. The protein encoded by this gene is one of the PKC family members. This protein kinase is expressed solely in the brain and spinal cord and its localization is restricted to neurons. It has been demonstrated that several neuronal functions, including long term potentiation (LTP) and long term depression (LTD), specifically require this kinase. Knockout studies in mice also suggest that this kinase may be involved in neuropathic pain development. Defects in this protein have been associated with neurodegenerative disorder spinocerebellar ataxia-14 (SCA14).

Storage:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. Avoid multiple freeze-thaw cycles.

To place an order, please Click HERE.





