

CD3e Human

Description: CD3e Human Recombinant produced in E. coli is a single polypeptide chain containing 127 amino acids (23-126) and having a molecular mass of 14.1kDa. CD3e is fused to a 23 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #: PRPS-1324

Synonyms: T-cell surface glycoprotein CD3 epsilon chain, T-cell surface antigen T3/Leu-4 epsilon chain, CD3e, T3E, TCRE.

For research use only.

Source: E.coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MGSDGNEEMG GITQTPYKVS
ISGTTVILTC PQYPGSEILW QHNDKNIGGD EDDKNIGSDE DHLSLKEFSE LEQSGYYVCY
PRGSKPEDAN FYLYLRARVC ENCMEMD.

Purity: Greater than 90% as determined by SDS-PAGE.

Formulation:

The CD3e solution (1mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 0.2M NaCl and 10% glycerol.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

CD3-epsilon polypeptide (CD3E) along with CD3-gamma, -delta and -zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, creates the T-cell receptor-CD3 complex. This complex has an imperative role in coupling antigen recognition to a number of intracellular signal-transduction pathways. The genes encoding the epsilon, gamma delta polypeptides are found in the same cluster on chromosome 11. The epsilon polypeptide plays a vital role in T-cell development. CD3E gene defects cause immunodeficiency. CD3E is also linked to a susceptibility to type I diabetes in women.

To place an order, please [Click HERE](#).