www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

Phospho-STMN1-S38

Reactivity: Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:100

Calculated MW:19kDa

Observed MW:Refer to Figures

Immunogen:

A phospho specific peptide corresponding to residues surrounding S38 of human STMN1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Concentration:

b

Synonym:

Lag; SMN; OP18; PP17; PP19; PR22; LAP18; C1orf215;

Background:

This gene belongs to the stathmin family of genes. It encodes a ubiquitous cytosolic phosphoprotein proposed to function as an intracellular relay integrating regulatory signals of the cellular environment. The encoded protein is involved in the regulation of the microtubule filament system by destabilizing microtubules. It prevents assembly and promotes disassembly of microtubules. Multiple transcript variants encoding different isoforms have been found for this

To place an order, please Click HERE.



Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:3925

Isotype:IgG

Swiss Prot:P16949

Purity: Affinity purification

For research use only.





