

Leptin Mouse

Description: Leptin Mouse Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 147 amino acids and having a molecular mass of 16.2 kDa. The Leptin is purified by proprietary chromatographic techniques.

Synonyms: OB Protein, Obesity Protein, OBS, Obesity factor.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: MVPIQKVQDD TKTLIKTIVT RINDISHTQS VSAKQRTVGL DFIPGLHPIL
SLSKMDQTLA VYQQVLTSLP SQNVLQIAND LENLRDLLHL LAFSKSCSLP QTSG LQKPES
LDGVLEASLY STEVVALSRL QGSLQDILQQ LDVSPEC.

Purity: Greater than 95.0% as determined by: (a) Analysis by SEC-HPLC. (b) Analysis by SDS-PAGE.

Formulation:

The mouse Leptin was lyophilized from a concentrated (1mg/ml) solution containing 0.1% TFA.

Stability:

Lyophilized Leptin although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Leptin should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Leptin in sterile 18M-cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Introduction:

A 16-kDa peptide hormone secreted from white adipocytes and implicated in the regulation of food intake and energy balance. Leptin provides the key afferent signal from fat cells in the feedback system that controls body fat stores.

Biological Activity:

Biological activity of Mouse Leptin is performed in two different mouse obesity models, ob/ob and NZO. Both strains of mice were treated via intraperitoneal injection once daily at a dose of 5µg Leptin/gram body weight for a period of 14 days. Significant effects on body weight, food consumption, and plasma glucose levels were observed to saline-treated controls.

References:

Title: Severe pulmonary metastasis in obese and diabetic mice. Publication: Article first published online: 22 SEP 2006 DOI: 10.1002/ijc.22248 Copyright

Catalog #:CYPs-358

For research use only.