www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

IFN a 2a Human, Plant

Description:Interferon Alpha Human 2a Recombinant produced in Plant is a single, glycosylated, polypeptide chain containing 165 amino acids and having a molecular mass of 19 kDa. The Interferon-a 2a contains affinity 6xHis tag on C-terminus. The IFN-A 2a is purified by proprietary chromatographic techniques.

Catalog #:CYPS-550

For research use only.

Synonyms:Leukocyte interferon, B cell interferon, Type I interferon, IFNA2, IFN-2a.

Source: Nicotiana Sp. Plant.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: The sequence of the first twelve N-terminal amino acids was determined and was found to be Cys-Asp-Leu-Pro-Gln-Thr-His-Ser-Leu-Gly-Ser-Arg.

Purity: Greater than 99.0% as determined by both(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Formulation:

Lyophilized, containing 50% SDS.

Stability:

Lyophilized Interferon alpha 2a although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IFN-alpha 2a should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Interferon-alpha 2a in sterile 18M-cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Introduction:

IFN-alpha is produced by macrophages and has antiviral activities. Interferon stimulates the production of two enzymes: protein kinase and an oligoadenylate synthetase.

Biological Activity:

The specific activity was determined by a quantitative gene report bioassay using human Type I interferon-sensitive cells. Compared with bacterial derived recombinant IFN-a 2a, assayed in the same conditions, plant derived IFN-alpha 2a showed 6.1

To place an order, please Click HERE.





