

Leptin tA Mouse, PEG

Description: Leptin Antagonist Triple Mutant Mouse Recombinant is a single non-glycosylated polypeptide chain containing 146 amino acids and additional Ala at N-terminus and having a molecular mass of ~ 16 kDa. The Mouse Leptin antagonist was mutated, resulting in L39A/D40A/F41A mutant. The Mouse Leptin antagonist is bound to 20 kDa mono-PEG at N-terminus, resulting in 35.6 kDa. The Mouse Leptin triple antagonist runs as a 48 kDa. Leptin Antagonist Triple Mutant Mouse Recombinant was purified by proprietary chromatographic techniques.

Catalog #: CYP5-573

For research use only.

Source: Escherichia coli.

Physical Appearance: White lyophilized (freeze-dried) powder.

Purity: Greater than 99.0% as determined by: (a) Gel filtration analysis. (b) Analysis by SDS-PAGE.

Formulation:

The Mouse Leptin triple antagonist was lyophilized from a concentrated (0.65mg/ml) solution with 0.003M NaHCO₃.

Stability:

Lyophilized Leptin Antagonist Triple Mutant Mouse Recombinant although stable at room temperature for several weeks, should be stored desiccated below -18°C. Upon reconstitution at > 0.1 Leptin mutant mg/ml and up to 2mM and filter sterilization LEP mutant can be stored at 4°C or even room temperature for several weeks making it suitable for long term infusion studies using osmotic pumps. At lower concentration addition of a carrier protein (0.1% HSA or BSA) is suggested. Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Leptin Antagonist Triple Mutant Mouse Recombinant in sterile water or sterile 0.4% NaHCO₃ adjusted to pH 8-9, not less than 100

Biological Activity:

Leptin Antagonist Triple Mutant Mouse Recombinant half-life in circulation after SC injection was over 20 hours. Leptin Antagonist Triple Mutant Mouse Recombinant is capable of inhibiting leptin-induced proliferation of BAF/3 cells stably transfected with the long form of human leptin receptor. Leptin Antagonist Triple Mutant Mouse Recombinant in vitro activity is 5-6 fold lower than the non-pegylated antagonist, though in vivo it has profound weight gain effect (as compared to the non-pegylated antagonist), resulting mainly from increased food intake.

To place an order, please [Click HERE](#).