

## TSLP Human

**Description:** TSLP Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 132 amino acids and having a molecular mass of 15 kDa. The TSLP is purified by proprietary chromatographic techniques.

**Synonyms:** Thymic Stromal Lymphopoietin, TSLP.

**Source:** Escherichia Coli.

**Physical Appearance:** Sterile Filtered White lyophilized (freeze-dried) powder.

**Amino Acid Sequence:** MYDFTNCDFE KIKAAAYLSTI SKDLITYMSG TKSTEFNNTV  
SCSNRPHCLT EIQLTFNPT AGCASLAKEM FAMKTKAALA IWCPGYSETQ INATQAMKKR  
RKRKVTTNKC LEQVSQLQGL WRRFNRPLLK QQ.

**Purity:** Greater than 97.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

**Formulation:**

Lyophilized from a concentrated solution (1mg/ml) contains 130mM NaCl, and 20mM sodium phosphate, pH 7.4.

**Stability:**

Lyophilized TSLP although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution TSLP should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

**Usage:**

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Solubility:**

It is recommended to reconstitute the lyophilized TSLP in sterile 18M-cm H<sub>2</sub>O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.


**Introduction:**

TSLP protein is a hemopoietic cytokine which signals throughout a heterodimeric receptor complex composed of the thymic stromal lymphopoietin receptor & the Interleukin-7 receptor alpha chain. TSLP impacts myeloid cells thus induces the discharge of T cell-attracting chemokines from monocytes & increases the growth of CD11c(+) dendritic cells. TSLP is mainly expressed in the heart, liver and prostate. TSLP is related in its biological activities with IL-7 and binds with the heterodimeric receptor complex consisting of the Interleukin-7 receptor alpha chain & the TSLPR. Similar to IL-7, TSLP enhances phosphorylation of STAT3 and STAT5, though uses kinases excluding JAKs for its activation. TSLP induces the release of T cell-attracting chemokines such as TARC & MDC from monocytes & triggers CD11c(+) dendritic cells. TSLP activated dendritic cells primes naive T cells to manufacture pro-allergic cytokines such as Interleukin-4, Interleukin-5, Interleukin-13 and TNF-alpha whereas down-regulating Interleukin-10 and IFN-gamma play a role in the initiation of allergic inflammation.

**Biological Activity:**

The ED50 is determined by the ability to induce proliferation of human Interleukin-7 receptor alpha & human TSLP Receptor co-transfected with mouse BaF3 cells which was found to be less than 0.3ng/ml.

**To place an order, please [Click HERE](#).**



Catalog #:CYP5-579

For research use only.