

Leptin tA Ovine, PEG

Description: Pegylated Leptin Antagonist Triple Mutant Ovine Recombinant is a single non-glycosylated polypeptide chain containing 146 amino and additional Ala at N-terminus acids and having a molecular mass of ~ 35.6 kDa, Leptin was mutated, resulting in L39A/D40A/F41A mutant. However due to enlarged hydrodynamic volume it runs on the SDS-Page as 48 kDa protein and in gel-filtration on Superdex 200 as over 200kDa protein. Leptin Antagonist Triple Mutant Ovine Recombinant Mono-Pegylated with 20kDa PEG and was purified by proprietary chromatographic techniques.

Catalog #: CYPs-710

For research use only.

Source: Escherichia coli.

Physical Appearance: White lyophilized (freeze-dried) powder.

Purity: Greater than 95.0% as determined by: (a) Gel filtration analysis. (b) Analysis by SDS-PAGE.

Formulation:

The protein was lyophilized from a concentrated (0.65mg/ml) solution with 0.003mM NaHCO₃.

Stability:

Lyophilized Leptin Antagonist Triple Mutant Ovine Recombinant although stable at room temperature for several weeks, should be stored desiccated below -18C. Upon reconstitution at > 0.1 Lep-tA mutant mg/ml and up to 2mM and filter sterilization Leptin mutant can be stored at 4C or even room temperature for several weeks making it suitable for long term infusion studies using osmotic pumps. At lower concentration addition of a carrier protein (0.1% HSA or BSA) is suggested. Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Leptin-Antagonist Triple Mutant Ovine Recombinant in sterile water or sterile 0.4% NaHCO₃ adjusted to pH 8, not less than 100

Biological Activity:

Pegylated triple antagonist is capable of inhibiting leptin-induced proliferation of BAF/3 cells stably transfected with the long form of human leptin receptor. Its in vitro activity is 6-8 fold lower than the non-pegylated antagonist but in vivo it has profound weight gain effect (as compared to the non-pegylated antagonist like in mouse leptin antagonists), resulting mainly from increased food intake.

To place an order, please [Click HERE](#).