www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

# MMP 8 Human

Description: Matrix Metalloproteinase-8 Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain having a molecular mass of 75 kDa. The MMP-8 is purified by proprietary chromatographic techniques.

For research use only.

Catalog #:ENPS-308

Synonyms: Neutrophil collagenase, EC 3.4.24.34, Matrix metalloproteinase-8, MMP-8, PMNL collagenase, PMNL-CL, HNC, CLG1.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered clear solution.

Purity: Greater than 90% as determined by SDS-PAGE.

#### Formulation:

The MMP-8 protein solution (100 units/ml) in 0.05M Tris-HCl buffer, pH 7.6, 0.2M NaCl, 5mM CaCl2, 0.0025% NaN3 and 0.1% BSA.

#### Stability:

MMP-8 although stable at  $4^{\circ}\text{C}$  for 1 week, should be stored desiccated below -18°C. Please prevent freeze-thaw cycles.

#### Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

## Applications:

Used as a standard for analyzing mammalian collagenase activity.

### Introduction:

Full-length recombinant human neutrophil pro-collagenase (MMP-8), latent form. Matrix metalloproteinase 8 (MMP-8), or neutrophil collagenase, degrades interstitial collagens, acting preferentially on collagen type I. Increased full-length MMP-8 protein was associated with infiltration into the skin of neutrophils, which are the major cell type that expresses MMP-8.MMP-8 is synthesized and stored in specific granules in neutrophil leukocytes. MMP-8 activity is therefore regulated by factors such as surface-bound ligands (IgG or complement components) that release it through degranulation. Once released and activated through proteolytic or oxidative mechanisms, MMP-8 plays a major role in the connective tissue turnover that accompanies inflammatory processes.

# **Biological Activity:**

100 units/ml after activation with APMA by solution assay method. One unit of collagenolytic activity is defined as the cleavage of 1

To place an order, please Click HERE.





