

PRDX6 Human

Description: Peroxiredoxin-6 Human Recombinant fused with a 20 amino acid His tag at N-terminus produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 244 amino acids (1-224 a.a.) and having a molecular mass of 27.1kDa. The Peroxiredoxin-6 is purified by proprietary chromatographic techniques.

Synonyms: Peroxiredoxin-6, Antioxidant protein 2, 1-Cys peroxiredoxin, Acidic calcium-independent phospholipase A2, Non-selenium glutathione peroxidase, 24 kDa protein, Liver 2D page spot 40, Red blood cells page spot 12, 1-Cys PRX, aiPLA2, NSGPx, PRDX6, AOP2, KIAA

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSSLVPRGSH MPGGLLGLDV APNFEANTTV
GRIRFHDFLG DSWGILFSP RDFTPVCTTE LGRAAKLAPE FAKRNVKLIASIDSVEDHL
AWSKDINAYN CEEPTEKLPF PIIDDRNREL AILLGMLDPA EKDEKGMPTV ARVVFVFGPD
KKLKSILYP ATTGRNFDEI LRVVISLQLT AEKRVATPVD WKDGDSVMVL PTIPEEEAKK
LFPKGVFTKE LP

Purity: Greater than 95.0% as determined by SDS-PAGE.

Formulation:

The Peroxiredoxin-6 solution contains 20mM Tris-HCl buffer (pH8.0) and 20% Glycerol.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Peroxiredoxin 6 (PRDX6) belongs to the thiol-specific antioxidant protein family. PRDX6 is a bifunctional enzyme with 2 distinct active sites. PRDX6 is involved in redox regulation of the cell and can reduce Hydrogen peroxide and short chain organic, fatty acid, and phospholipid hydroperoxides. PRDX6 may have a role in the regulation of phospholipid turnover as well as in protection against oxidative injury. Furthermore, PRDX6 eases the oxidative stress and TGF-beta-induced abnormalities of human trabecular meshwork cells. In addition, PRDX6 is necessary for blood vessel integrity in injured skin. At acidic pH, PRDX6 binds to reduced phospholipids, however at cytosolic pH PRDX6 binds only to phospholipids that are oxidized which is compatible with the role for PRDX6 in the repair of peroxidized cell membranes. Hydrogen peroxide-mediated hyperoxidation of PRDX6 induces cell cycle arrest at the G2/M transition via up-regulation of iPLA2 activity. Overexpression of PRDX6 is linked to oligodendroglioma.

Biological Activity:

www.neobiolab.com

info@neobiolab.com

888.754.5670, +1 617.500.7103 United States

0800.088.5164, +44 020.8123.1558 United Kingdom

The specific activity was found to be approximately 95-120 pmole/min/



To place an order, please [Click HERE](#).

Catalog #:ENPS-439

For research use only.

© 2015 NeoScientific



neobiolab | [Term](#) | [Quote/Order](#) | [Services](#) | [Products](#) | [Support](#) | [Corporate](#) | [Contact Us](#)

Copyrights @ 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)



Request a
CATALOG
[CLICK HERE](#)