

## FKBP2 Human

**Description:**FKBP2 Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 122 amino acids (22-142 a.a.) and having a molecular mass of 13.4 kDa. The FKBP2 is purified by proprietary chromatographic techniques.

**Catalog #:**ENPS-510

For research use only.

**Synonyms:**FKBP13, FKBP-2, Peptidyl-prolyl cis-trans isomerase FKBP2, PPIase FKBP2, FK506-binding protein 2, Rotamase, Immunophilin FKBP13, 13 kDa FK506-binding protein, 13 kDa FKBP, FKBP-13, FKBP2, PPIase.

**Source:**Escherichia Coli.

**Physical Appearance:**Sterile Filtered clear colorless solution.

**Amino Acid Sequence:**MATGAEGKRK LQIGVKKRVD HCPIKSRKGD VLHMHYTGKL  
EDGTEFDSSL PQNQPFVFSL GTGQVIKWD QLLGMCEGE KRKLVIPSEL GYGERGAPPK  
IPGGATLVFE VELLKIERRT EL.

**Purity:**Greater than 90% as determined by SDS-PAGE.

**Formulation:**

The FKBP2 solution contains 20mM Tris-HCl pH-8, 1mM DTT, and 10% glycerol.

**Stability:**

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

**Usage:**

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Introduction:**

FKBP2 is part of the immunophilin protein family, which takes part in immunoregulation and basic cellular processes involving protein folding and trafficking. FKBP2 is a cis-trans prolyl isomerase that binds the immunosuppressants FK506 and rapamycin. FKBP2 functions as an ER chaperone and as a component of membrane cytoskeletal scaffolds.

**Biological Activity:**

Specific activity is > 270 nmoles/min/μg, and is defined as the amount of enzyme that cleaves 1 μmole of suc-AAFP-pNA per minute at 25°C in Tris-HCl pH8.0 using chymotrypsin.

**To place an order, please [Click HERE](#).**