www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

HAAO Human

Description: HAAO Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 310 amino acids (1-286 a.a.) and having a molecular mass of 35kDa.HAAO is fused to a 24 amino acid His-tag at N-terminus & Durified by proprietary chromatographic techniques.

Catalog #:ENPS-624

For research use only.

Synonyms: 3-hydroxyanthranilate 3,4-dioxygenase, 3-hydroxyanthranilate oxygenase, 3-HAO, 3-hydroxyanthranilic acid dioxygenase, HAD, HAAO, HAO.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MGSHMERRLG VRAWVKENRG SFQPPVCNKL MHQEQLKVMF IGGPNTRKDY HIEEGEEVFY QLEGDMVLRV LEQGKHRDVV IRQGEIFLLP ARVPHSPQRF ANTVGLVVER RRLETELDGL RYYVGDTMDV LFEKWFYCKD LGTQLAPIIQ EFFSSEQYRT GKPIPDQLLK EPPFPLSTRS IMEPMSLDAW LDSHHRELQA GTPLSLEGDT YE

Purity: Greater than 90.0% as determined by SDS-PAGE.

Formulation:

HAAO protein solution (0.5mg/ml) containing 20mM Tris-HCl buffer (pH 8.0), 0.15M NaCl, 10% glycerol and 1mM DTT.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

HAAO is a monomeric cytosolic protein which is a member of intramolecular dioxygenases family containing nonheme ferrous iron. The HAAO protein catalyzes the synthesis of quinolinic acid (QUIN) from 3-hydroxyanthranilic acid. QUIN is an excitotoxin whose toxicity is facilitated by its ability to activate glutamate N-methyl-D-aspartate receptors. Amplified cerebral levels of QUIN may contribute to the pathogenesis of neurologic and inflammatory disorders. HAAO is widely distributed in peripheral organs, such as liver and kidney, and is also present in low amounts in the central nervous system.

To place an order, please Click HERE.





