

AMD1 Human

Description:AMD1 Human Recombinant produced in E. coli is a single polypeptide chain containing 292 amino acids (68-334) and having a molecular mass of 33.4 kDa.AMD1 is fused to a 25 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #:ENPS-631

For research use only.

Synonyms:Adenosylmethionine decarboxylase 1, S-adenosylmethionine decarboxylase proenzyme, AdoMetDC, S-adenosylmethionine decarboxylase 1, SAMDC, AMD, EC 4.1.1.50.

Source:E.coli.

Physical Appearance:Sterile Filtered colorless solution.

Amino Acid Sequence:GSSHHHHHH SSGLVPRGSH MGSMMSSMFV SKRRFILKTC
GTTLLKALV PLLKLARDYS GFDSIQSFFY SRKNFMKPSH QGYPHRNFE EIEFLNAIFP
NGAAYCMGRM NSDCWLYLTL DFPESRVISQ PDQTLILMS ELDPAVMDQF YMKDGVTAKD
VTRESGIRDL IPGSVIDATM FNPCGYSMNG MKSDGTYWTI HITPEPEFSY VSFETNLSQT
SYDDLIRKVV EVF

Purity:Greater than 80% as determined by SDS-PAGE.

Formulation:

The AMD1 solution (0.5mg/1ml) contains 20mM Tris-HCl buffer (pH 8.0), 100mM NaCl, 1mM DTT and 20% glycerol.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Adenosylmethionine decarboxylase proenzyme (AMD1) is synthesized originally as an inactive proenzyme. Putrescine stimulates both the proenzyme processing and the catalytic activity. The catalytic activity is inhibited by iodoacetic acid. The active enzyme formation entails a self-maturation process in which the active site pyruvoyl group is produced from an internal serine residue using an autocatalytic post-translational modification.

To place an order, please [Click HERE](#).