

HAVCR2 Human

Description: Recombinant Human HAVCR2 produced in E. coli is a single polypeptide chain containing 206 amino acids (aa 22-202) and having a molecular mass of 22.7kDa. HAVCR2 is fused to a 25 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #: HAPS-231

For research use only.

Synonyms: Hepatitis A virus cellular receptor 2, HAVcr-2, T-cell immunoglobulin and mucin domain-containing protein 3, TIMD-3, T-cell membrane protein 3, TIM-3, HAVCR2, TIM3, TIMD3, TIM3, KIM-3.

Source: Escherichia Coli.

Physical Appearance: Sterile filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MGSHMEPGAG HLDGHRAGSP
SLRQALCDGS AVMFSSKERG RCTVINFLVPL EAPLRSTPRS RQVTEACGGE GRAVPLGSEP
EWSVGMEAT LEQHLEDTKM NPSIVGVLC TDSQGLNLGCR GTLSDEHAGV ISVLAQQAQK
LTSDPTDIPV VCLESNDNGNI MIQKHDGIV AVHKMAS.

Purity: Greater than 90% as determined by SDS-PAGE.

Formulation:

The HAVCR2 solution (1mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 0.4M urea and 10% glycerol.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Hepatitis A Virus Cellular Receptor 2 (HAVCR2) is a member of the immunoglobulin superfamily. HAVCR2 controls macrophage activation. HAVCR2 inhibits T-helper type 1 lymphocyte (Th1)-mediated auto- and alloimmune responses and stimulates immunological tolerance.

To place an order, please [Click HERE](#).