

FLT1 Human

Description: Soluble FLT1 Human Recombinant produced in baculovirus is monomeric, glycosylated, polypeptide containing 688 amino acids and having a molecular mass of 96 kDa. The soluble receptor protein contains only the first 6 extracellular domains, which contain all the information necessary for binding of VEGF. The FLT1 is purified by proprietary chromatographic techniques.

Synonyms: FLT-1, FLT1, Tyrosine-protein kinase receptor FLT, Flt-1, Tyrosine-protein kinase FRT, Fms-like tyrosine kinase 1, VEGFR-1.

Source: Insect Cells.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: MVS YWDTGVL LCALLSCLLL TGSSSGSKLK DPELSLKGQTQ
HIMQAGQTLH LQCRGEAAHK WSLPEMVSKE SERLSITKSA CGRNGKQFCS TLTNTAQAN
HTGFYSCKYL AVPTSKKKET ESAIYIFISD TGRPFVEMYS EIPEIIHMTE GRELVIPCRV
TSPNITVTLK KFPLDTLIPD GKRIIWDSRK GFISNATYK EIGLLTCEAT VNGHLYKTNV
LTHRQNTNII DV

Purity: Greater than 90.0% as determined by (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Formulation:

FLT1 was lyophilized from a concentrated (1 mg/ml) sterile solution containing 1x PBS.

Stability:

Lyophilized FLT-1 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution FLT1 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized FLT1 in sterile water not less than 100

Introduction:

Endothelial cells express three different vascular endothelial growth factor (VEGF) receptors, belonging to the family of receptor tyrosine kinases (RTKs). They are named VEGFR-1 (Flt-1), VEGFR-2 (KDR/Flk-1), VEGFR-3 (Flt-4). Their expression is almost exclusively restricted to endothelial cells, but VEGFR-1 can also be found on monocytes, dendritic cells and on trophoblast cells. The flt-1 gene was first described in 1990. The receptor contains seven immunoglobulin-like extracellular domains, a single transmembrane region and an intracellular split tyrosine kinase domain. Compared to VEGFR-2 the Flt-1 receptor has a higher affinity for VEGF but a weaker signaling activity. VEGFR-1 thus leads not to proliferation of endothelial cells, but mediates signals for differentiation. Interestingly a naturally occurring soluble variant of VEGFR-1 (sVEGFR-1) was found in HUVE supernatants in 1996, which is generated by alternative splicing of the flt-1 mRNA.

Biological Activity:

The activity of FLT1 was determined by its ability to abolish the binding of iodinated VEGF to solid surfaces or cell surface receptors. The ED50 for this effect is typically 10 ng/ml, corresponding to a specific activity of 100,000IU/mg.

Catalog #:PKPS-247

For research use only.

To place an order, please [Click HERE](#).