www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

# Trypsin Bovine

Description: Recombinant Bovine Trypsin is free from any animal and human sources.% of inhibition is equal to bovine trypsin.

Catalog #:PRPS-320

Source: Corn.

For research use only.

Physical Appearance: Sterile Filtered lyophilized powder.

Purity: Greater than 90% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

#### Formulation:

The protein was lyophilized without any additives.

#### Stability:

Bovine Trypsin although stable at room temp for 1 week, should be stored desiccated below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

## Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

## Solubility:

It is recommended to reconstitute the lyophilized Bovine Trypsin in sterile 18M-cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

## Introduction:

Trypsin is a serine protease that hydrolyses proteins, it is found in the digestive system of numerous vertebrates. Trypsin is produced as the inactive proenzyme trypsinogen in the pancreas. Trypsin cleaves peptide chains at the carboxyl side of the amino acids lysine and arginine, except when either is followed by proline. Trypsin is secreted into the duodenum, where it acts to hydrolyses peptides into amino acids, which is necessary for the uptake of protein in the food even though peptides are smaller than proteins; they are still too big to be absorbed through the lining of the ileum. The optimal operating pH for Trypsins is about 8 and about 37°C temperature. In cystic fibrosis disease there is a deficiency in transport of trypsin and other digestive enzymes from the pancreas. Trypsin is widely used in various biotechnological processes since its available in high quantity in the pancreases, and can be purified rather easily.

# **Biological Activity:**

3,650 Units/mg.

To place an order, please Click HERE.





