

## ANXA1 Human

**Description:**ANXA1 produced in E.Coli is a single, non-glycosylated polypeptide chain containing 346 amino acids (1-346 a.a.) and having a molecular mass of 38.7 kDa. ANXA1 is purified by proprietary chromatographic techniques.

**Catalog #:**PRPS-686

For research use only.

**Synonyms:**ANX1, LPC1, ANXA1, Lipocortin I, Calpactin II, Chrombindin-9, p35, Annexin-1, Phospholipase A2 inhibitory protein, Annexin I, Annexin A1.

**Source:**Escherichia Coli.

**Physical Appearance:**Sterile filtered colorless solution.

**Amino Acid Sequence:**MAMVSEFLKQ AWFIEEEQE YVQTVKSSKG GPGSAVSPYP  
TFNPSSDVAA LHKAIMVKGV DEATIIDILT KRNNAGRQKI KAAYLQETGKPLDETLKKAL  
TGHLEEVVLA LLKTPAQFDA DELRAAMKGL GTDEDTLIEI LASRTNKEIR DINRVYREEL  
KRDIAKDITS DTSGDFRNAL LSLAKGDRSE DFGVNEDLAD SDARALYEAG ERRKGTDVNV  
FNTILTTRSY PQL

**Purity:**Greater than 90.0% as determined by SDS-PAGE.

### Formulation:

The ANXA1 protein solution contains 20mM Tris-HCl, pH-8, 100mM NaCl, 1mM DTT and 10% glycerol.

### Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

### Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

### Introduction:

ANXA1 is part of the family of Ca(2+)-dependent phospholipid binding proteins which have a Mw between 35kDa-40kDa and are situated on the cytosolic face of the plasma membrane. ANXA1 protein has a Mw of 40kDa, with phospholipase A2 inhibitory activity to bind from two to four calcium ions with high affinity. Since phospholipase A2 is necessary for the biosynthesis of the potent mediators of inflammation, prostaglandins and leukotrienes, ANXA1 might have potential anti-inflammatory activity. ANXA1 promotes membrane fusion and plays a role in exocytosis. The recognition of ANXA1 protein by immunocytochemical leads a simple, highly sensitive and specific assay for diagnosis of hairy cell leukemia.

**To place an order, please [Click HERE](#).**